

HPV – THE MOST COMMON SEXUALLY TRANSMITTED VIRUS

Human papillomavirus (HPV) is a common infection that affects skin and mucous membranes. Some types cause common skin warts in areas such as the hands or feet. Some types cause warts in the genital areas. Certain high-risk types can cause cancers of the cervix, vagina, vulva, penis, anus, and throat. More than 150 viral types of HPV have been identified – about 40 of these are associated with sexually transmitted infections, and people may be infected with more than one type at a time (NCI, 2011; Gillison et al., 2012; Schiffman & Castle, 2003; Wiley et al., 2002). HPV has affected humans for thousands of years – ancient Greek and Roman medical records described genital lesions consistent with genital warts and associated them with sexual activity (Jay & Moscicki, 2000).

Today HPV is the most common sexually transmitted infection in the U.S. – yet in 2013, 35 percent of Americans had never heard of it and 89 percent had never discussed HPV with their health care provider (NCI, 2013). About 79 million Americans are currently infected with HPV, and approximately 14 million Americans acquire HPV annually. As many as half of these infections are among young people between the ages of 15 and 24 (CDC, 2014a). About one in four of all young adults in America have HPV at any given time (Dunne et al., 2007; Weinstock et al., 2004). HPV is also prevalent among people with immunosuppressive disorders, such as HIV (Koutsky & Kiviat, 1999). HPV is so common, in fact, that it is considered a virtual marker for having had sex (Boonstra, 2004). In fact, most sexually-active people will get at least one type of HPV at some point in their lives (CDC, 2014a).

Transmission

HPV is transmitted by direct skin-to-skin contact with an infected individual. Transmission is usually from vaginal, oral, or anal sexual contact and can occur whether or not warts or other symptoms are present (McDermott-Webster, 1999). Unprotected penetrative intercourse with multiple partners is the greatest behavioral risk for contraction of HPV (Kjaer et al., 2001; Winer et al., 2003).

The virus can also be transmitted from mother to infant during childbirth – also known as vertical transmission (Puranen, 1997). In one of the largest studies to look at both oral and genital HPV infections in newborns, research showed that the vertical transmission rate was less than one percent (Smith et al., 2004). The development of recurrent respiratory papillomatosis (warts in the respiratory tract) is one potential consequence of vertical transmission (Smith et al., 2004; Kashima et al., 1996). It is estimated that about 2,000 out of every four million newborns are infected (Jay & Moscicki, 2000). This is a serious, and potentially fatal, condition that may require frequent laser surgery to prevent obstruction of the infant's airways (NIAID, 2004).

Some research also suggests that genital HPV can be transmitted through nonsexual routes, via fomites – inanimate objects such as towels or underwear – but more research must be conducted to examine these modes of transmission (Carson, 1997; Keller et al., 1995; Stevens-Simon et al., 2000). According to the CDC, transmission by fomites has never been documented (CDC, 2013).

Natural History

Although there is currently no “cure” for genital HPV infection, most cases are transient and clear themselves without medical intervention (Brown et al., 2005; CDC, 2014a; Elfgren et al., 2000; Ho et al., 1998). It is estimated that approximately 90 percent of HPV infections are cleared by the body’s immune system within two years (CDC, 2014c). The viral type of HPV is a major determinant in the duration of infection, with types 16, AE7, 61, 18, and 73 having the longest average duration (Elfgren et al., 2000; Ho et al., 1998; Muñoz et al., 2004; Richardson et al., 2003; Woodman et al., 2001). Other factors associated with the persistence of HPV infections are age (older than 30 years), infection with multiple HPV types, and a compromised immune system (CDC, 2014a; Hildesheim et al., 1994; Ho et al., 1995; Moscicki et al., 1998).

Subclinical Manifestations of HPV

Most HPV infections are subclinical (no visible signs or symptoms), and many people with HPV never know they have it (Verdon, 1997). HPV targets the deep, basal level of the skin and most often causes no clinical or microscopic changes in the cells of the skin (Keller et al., 1995; Verdon, 1997). In some cases, subclinical HPV may cause cellular changes that are only detectable using clinical instruments or the study of cervical cells. These changes may be, in rare instances, the precursor to cancer cells (Lytwyn & Sellors, 1997).

Changes related to HPV infections that cannot be seen with the naked eye may be identified using a variety of clinical tools:

- A hand lens or colposcope may be used to magnify cervical and vaginal tissue (Verdon, 1997).
- Pap tests may reveal precancerous conditions of the cervix that are caused by HPV. (Some experts also recommend anal Pap tests for men at increased risk of anal cancer – men who have sex with men and men who are HIV positive) (CDC, 2014a; Gilden, 2005; Tuller, 2003).
- HPV testing of samples taken with a cervical swab can detect high-risk types of HPV.

In March 1999, the U.S. Food and Drug Administration (FDA) approved the Hybrid Capture II HPV test – a DNA-based technology that can detect 13 high-risk types of HPV (those associated with an increased risk of cancer) (“HPV DNA Tests”, 2000). Today, there are four HPV tests currently available for clinical use (NCCCC, 2014). Starting at age 25, women can receive the Roche HPV test in lieu of a Pap test every three years. HPV tests are not recommended along with Pap tests for women under the age of 30 unless they have atypical or unclear Pap test results. This is because HPV is very common; cervical cancer is rare at this age; and most HPV infections go away by themselves without causing any health problems. For women age 30 or older, HPV tests can be done at the same time as a Pap test. If both test results are normal, a woman has a very low risk of developing cervical cancer. She will not need a Pap and HPV test for five years. Some women age 30 or older see this choice as more appealing than having a Pap test more frequently (ACOG, 2009 and 2013).

Clinical Manifestations of HPV

In some instances HPV infection can lead to clinical manifestations that can be seen with the “naked” eye. Clinical manifestations can appear as classical warts, or as a variety of lesions on the cervix, vagina, vulva, penis, anus, or throat.

Genital Warts

Genital warts (condylomata acuminata) are the most common clinical manifestation of genital HPV. In more than 90 percent of cases, they are caused by HPV types 6 and 11, which are considered low-risk types because they are not associated with increased risk of cancer (Jay & Moscicki, 2000; Moscicki, 2005; Wiley, 2002).

It is estimated that one percent of the sexually active American population has genital warts, and women and men have similar rates of infection (CDC, 2013; Cockerell, 1995; Jay & Moscicki, 2000; Moscicki, 2005). About 360,000 people in the United States get genital warts each year (CDC 2014a).

Genital warts usually start as small bumps that appear in the anogenital area. They may be single or clusters and have a cauliflower-like appearance as they grow larger. In women, genital warts may appear on the vulva, in the

vagina, on the cervix, groin, or in the anal area. In men, they appear on the foreskin, head or shaft of the penis, groin, and in the anal area, urethra, and scrotum (ASHA, 2014; Cockerell, 1995). Rarely, warts may also develop in the mouth or throat of a person who has had sexual contact with an infected person (Koutsky & Kiviat, 1999).

Genital warts usually are painless, but they may cause itching or irritation (Cockerell, 1995). Genital warts are very contagious, with an estimated rate of infection between 60 and 75 percent from unprotected exposure (NIAID, 2004; Soper, 2002). The incubation period for genital warts is between two weeks and eight months, but it may last for years after exposure (Anic and Giuliano, 2012; ASHA, 2014).

Treating Genital Warts

Because there is no cure for HPV infections, the purpose of treatment is to control outbreaks of warts. Although genital warts often fade away by themselves, they sometimes need to be treated. There are a variety of options to treat warts, including several chemicals that can be applied directly to genital warts:

- biochloroacetic acid (BCA)
- trichloroacetic acid (TCA)
- podofilox
- imiquimod

BCA and TCA are chemicals that must be applied by a clinician. Podofilox and imiquimod are two treatments that can be prescribed for use at home. Podofilox is a self-applied cream or gel that destroys wart tissue. Imiquimod, also a self-applied cream, is an immune system modulator. It works by boosting the immune system to fight HPV infection. Some of these treatments can cause local discomfort, and some cannot be used during pregnancy (ASHA, 2014; Holmes et al., 2008).

A clinician can also remove genital warts with cryotherapy (freezing off), electrocautery therapy (burning off), laser therapy, or surgery (ASHA, 2014).

HPV and Cancer

It is estimated that in 2015 there will be about 12,900 new cases of invasive cervical cancer in the United States, which will result in about 4,100 deaths (ACS, 2015a). Worldwide, about 530,000 new cases are diagnosed each year. Cervical cancer is the fourth most common type of cancer among women worldwide and one of the leading causes of cancer-related mortality in women in the developing world. It was responsible for 266,000 deaths in 2012, about 87 percent of which occurred in developing countries (Ferlay et al., 2014). The median age of diagnosis for cervical cancer is 48 years (CDC, 2012a). Most women diagnosed with cervical cancer are younger than age 50 (ACS, 2015b).

Due largely to routine screening using Pap tests, the number of deaths attributed to cervical cancer in the United States dropped more than 50 percent over the last 30 years (ACS, 2015b). The five-year survival rate for patients diagnosed with localized disease is 91 percent. The overall five-year survival rate for all stages of cervical cancer is 68 percent (ACS, 2015a).

African Americans experience a disproportionate number of deaths from cervical cancer – due mainly to underscreening in this population. From 2007–2011, the death rate was 4.2 per 100,000 for African-American women, compared to 2.0 per 100,000 for non-Hispanic white women. Latinas and Native Americans also have cervical cancer death rates that are above average (ACS, 2015a).

Since the late 1800s, researchers have suspected that cervical cancer was sexually transmitted. Medical reports noted that nuns and virgins were not likely to have cervical cancer, and that women who were married to men who traveled a great deal or who had previous wives who died of cervical cancer were more likely to develop cervical cancer (“The Cervical Cancer Virus,” 1995). Today, 15–20 types of HPV have been classified as oncogenic, and the DHHS has added HPV to the list of cancer-causing agents (Janicek & Averette, 2001; Kay, 2005; Muñoz et al., 2003; Schiffman & Castle, 2003; Wiley et al., 2002). Large studies have found that HPV is present in more than 99 percent of cervical cancer tumors (Clifford et al., 2003; Walboomers et al., 1999). HPV 16 and 18 are responsible for about 70 percent of all

cervical cancers. Other HPV types are associated with the remaining 30 percent of cases (Bosch & deSanjosé, 2003; Clifford et al., 2003; Shah, 1997).

Most HPV infections never lead to the development of cervical cancer – even in the absence of medical intervention – and appropriate management of precancerous cervical lesions detected by Pap tests has greatly reduced the rate of invasive cervical cancer (Ho et al., 1998; NCI, 2014). Only one out of 1,000 women with HPV develops invasive cervical cancer (ACOG, 2000).

HPV appears to be necessary, but not sufficient, to the development of cervical cancer. Besides HPV type, researchers believe there are several cofactors that may contribute to the development of cervical cancer. These may include alcohol consumption, smoking, diet, family history, HIV infection, hormonal factors – including multiple pregnancies and the use of both oral contraceptives and DES, low socioeconomic status, the presence of other sexually transmitted infections, such as chlamydia and/or herpes simplex virus 2, and having an uncircumcised male partner (ACS, 2015b; Anttila et al., 2001; Moscicki, 2005; NCI, 2014).

Certain high risk HPV types are also now considered to be a cause of many cancers of the vagina, vulva, anus, penis, and throat. Although each of these cancers occurs less frequently than cervical cancer, taken together they equal more than the number of cases of cervical cancer in the U.S. (ACS, 2015a). The average age for diagnosis of these cancers is significantly later than for cervical cancer. The median age of diagnosis for vaginal cancer is 69 years and 66 years for vulvar cancer. Anal cancer is typically diagnosed at 60 years of age for women and 56 years for men, and the average age of diagnosis for cancer of the penis is 68 years. The mean age for diagnosis of HPV-related throat cancer is 61 among women and 58 among men (CDC, 2012a). As is the case with cervical cancer, HPV 16 and HPV 18 are most often associated with vaginal, vulvar, anal, and penile cancers (Chaturvedi et al., 2008; Eng & Butler, 1997). HPV is also associated with 20 percent of oropharyngeal (primarily the tongue and tonsils) cancers and 90 percent of skin cancers in immunocompromised patients (González et al., 2002; Ryerson et al., 2008). Men are three times more likely than women to develop throat cancers (Ryerson et al., 2008).

Vaccination

Over the last few years, there have been great strides in the development and testing of vaccines against HPV:

Gardasil®

Gardasil®, a prophylactic, quadrivalent (HPV types 6, 11, 16, and 18) vaccine manufactured by Merck and Co., Inc., was approved June 8, 2006, by the U.S. Food and Drug Administration. Given in three injections over a six-month period, clinical trials have shown this vaccine to be both safe and effective (Koutsky et al., 2002; Mao et al., 2006; Skjeldestad et al., 2005; Villa et al., 2005).

Follow-up studies (3.5 years after vaccination) have shown an effectiveness rate of 94 percent against persistent HPV 16 infection (Mao et al., 2006). These studies have also found a 100 percent effectiveness rate in preventing the development of high-grade (CIN 2-3), pre-cancerous cervical lesions related to HPV 16 and 18 – which cause 70 percent of all cervical cancers (Koutsky et al., 2002; Mao et al., 2006; Muñoz et al., 2010; Villa et al., 2005). Gardasil also protects against the two types of HPV that cause genital warts and protects against cancers of the anus in women and men (CDC 2014b) and researchers are optimistic that it protects against HPV-related throat cancers as well (Gillison, 2012).

Except for local irritation at the injection site, side effects in the study group were similar to the placebo group (Kahn, 2005). An Institute of Medicine (IOM) review of the scientific evidence on adverse events associated with vaccines covered by the National Vaccine Injury Compensation Program (VICP) found that “evidence favors acceptance of a causal relationship” between HPV vaccine and anaphylaxis, but the evidence is “not firm enough to be described as convincing” (IOM, 2011).

Cervarix™

Cervarix™, a three-dose bivalent (HPV types 16 and 18) prophylactic vaccine manufactured by GlaxoSmithKline (GSK) was approved in October, 2009 (GSK, 2009).

Studies showed the vaccine to be safe and 100 percent effective in preventing HPV type 16 and 18 infections, and nearly 100 percent immunogenic over a period of 4.5 years that have been measured so far (Harper

et al., 2004; Harper et al., 2006). No serious adverse events related to the vaccine were reported during these studies (Kahn, 2005). Preliminary results of phase III clinical trials showed detectable HPV 16 and 18 antibody levels six months following the completion of the vaccination series to be at least 16 to 26 times higher than antibody levels seen after natural infection (GSK, 2006; Schwarz et al., 2006). Cervarix also protects against cancer of the anus and is presumed to protect against HPV-related cancers of the throat (Gillison, 2012; Kreimer et al., 2011).

Gardasil® 9

On December 10, 2014, the FDA approved Gardasil® 9 for use in females ages 9 through 26 and males ages 9 through 15. It is approved for the prevention of cervical, vulvar, vaginal and anal cancers caused by HPV types 16, 18, 31, 33, 45, 52 and 58, and for the prevention of genital warts caused by HPV types 6 or 11. Gardasil 9 adds protection against five additional HPV types – 31, 33, 45, 52 and 58 – which cause approximately 20 percent of cervical cancers and are not covered by previously FDA-approved HPV vaccines. Gardasil 9 is manufactured by Merck and Co., Inc (FDA, 2014). Clinical trials found that anti-HPV 6/11/16/18 responses were non-inferior to those generated by the current Gardasil quadrivalent vaccine, with 97.1 percent efficacy against HPV types 31/33/45/52/58 related cervical/vulvar/vaginal disease (Serrano et al., 2014).

Social Acceptance of STI Vaccines for the Young

The development of these HPV vaccines has the enormous potential to improve the reproductive health and well-being of women and men. And because prophylactic HPV vaccines are only effective in individuals not currently infected by the virus, it will be important for the vaccine to be administered to women and men before they become sexually active.

In 2013, 28 percent of young women in the ninth grade, and 32 percent of young men in the ninth grade had had sexual intercourse (Kann et al., 2014). To reach these young people before they become sexually active, the FDA approved Gardasil for girls and women from nine to 26 years old, and the Center for Disease Control and Prevention's (CDC) Advisory Committee on Immunization Practices (ACIP) has recommended that the vaccine be

routinely administered to girls and boys at age 11 and 12 (CDC, 2014a; FDA, 2006; Markowitz et al., 2011). ACIP guidelines recommend "catch up" immunization of young women (ages 13–26 years) and young men (ages 13–21 and through 26 years for high-risk men) who have not been previously vaccinated. Initial testing was done mostly on women 15–26 years of age.

One study estimated that the vaccination of 12-year-old girls could potentially reduce the number of HPV 16- and 18-related cervical cancer cases by more than 95 percent. The vaccination of both young boys and girls could further reduce the number of cervical cancer cases by an additional three percent (Taira et al., 2004). Protecting men from genital warts could also potentially break the chain of transmission of genital warts (Bor, 2006).

The young target age of vaccination has led many researchers, health care providers, parents, and the patients themselves to consider the unique issues related to the acceptance of an STI, or more specifically an HPV vaccine. Some of the many potential barriers to vaccination include the belief that young children already receive too many vaccinations, concern that immunization might lead to risky sexual behavior, concern about vaccine safety, a reluctance to immunize against an STI, and a reluctance to talk with young children about STIs and sexuality (Kahn et al., 2005; Mays et al., 2004). These potential barriers were very similar to those observed upon the development and approval of hepatitis B vaccines – the first vaccines developed to reduce the risk of contracting a virus that can cause cancer (Rosenthal et al., 1995).

When individuals and parents learn about the connection between HPV and cervical cancer their reluctance to vaccinate dissipates. Surveys of young adults have shown HPV and general STI vaccination acceptance rates between 74 and 89 percent (Boehner et al., 2003; Kahn et al., 2003). Surveys of parents have shown HPV vaccine acceptance rates between 73 and 84 percent (Davis et al., 2004; Mays et al., 2004). Vaccinating young children opens the door to parent-child and provider-child communication. The HPV vaccine presents a unique opportunity to both introduce young children to sexuality education and educate them about healthy sexual expression.

Cost

Parents and providers are also concerned about the costs associated with the HPV vaccines. The retail cost of HPV vaccines is about \$130 a dose – \$390 for the full, three-shot series. Most insurance plans cover the cost. The Vaccines for Children (VFC) program may be able to help cover the cost for those who are uninsured or those whose insurance does not cover the cost. Children younger than 19 years old are VFC-eligible if they have no insurance, are eligible for Medicaid, or are American Indian or Alaskan Native (CDC, 2012b).

HPV-vaccine manufacturers also help women and men who cannot afford their vaccines. GlaxoSmithKline has its Vaccine Access Program (1-877-822-2911), which serves low-income women and men who are 19–25 and too old for the VFC program. Merck has its Vaccine Patient Assistance Program (1-800-293-3881), which helps women and men over 19 who do not have insurance or cannot afford to pay for vaccination (NCI, 2011).

While the potential cost of the vaccine is high, it is not comparable to the total cost of HPV and cervical cancer in the United States. In 2009, the total cost attributed to HPV in the United States was approximately \$1.7 billion (Owusu-Edusei et al., 2013). A cost-study analysis of women's health plan enrollees determined that the average per-woman cost associated with the screening and treatment of cervical HPV-related diseases was \$26,415 (Insinga et al., 2004). Vaccines will not eliminate the total cost of screening or treatment, but they do have the potential to significantly reduce it.

Ongoing Prevention

Vaccine boosters might be necessary. Until other vaccines are developed that will protect against all oncogenic HPV types, women will need to continue to practice safer sex and receive regular Pap tests.

Abstinence and lifelong monogamy will continue to be the most effective ways to avoid HPV infection entirely. Even if Gardasil and Cervarix are 100 percent effective, they only prevent HPV types that cause 70 percent of cervical cancers. Gardasil 9 only prevents HPV types that cause 90 percent of cervical cancers. Women will still need screening to protect themselves against cancer

caused by types not covered by a vaccine. For most sexually active women, the most important preventive measure to protect themselves from developing cervical cancer will continue to be regular Pap tests (Janicek & Averette, 2001). Avoiding skin-to-skin contact with someone with HPV is the most effective, but not always practical, strategy to prevent HPV infection. And although condoms may not entirely eliminate the risk of transmitting HPV, they are recommended for risk reduction (ASHA, 2014; Winer, et al., 2006). A study published in the *New England Journal of Medicine* showed that women whose partners used condoms consistently and correctly during vaginal intercourse over a period of eight months were 70 percent less likely to acquire a new HPV infection than women whose partners used condoms less than five percent of the time (Winer et al., 2006).

Because HPV may shed beyond the covered area, however, condoms do not provide as complete protection as they do for some other pathogens, such as HIV and gonorrhea (Stone et al., 1999). The claims of condom-use opponents who suggest that condom use leads to increased numbers of HPV infections are false and alarmist. Condom use cannot be blamed for the high prevalence of HPV infection or the incidence of cervical cancer among women in the U.S. In fact, two Dutch studies found that condom use promotes the regression of HPV lesions in women and men, as well as the clearance of HPV in women (Hogewoning et al., 2003; Bleeker et al., 2003).

While HPV is endemic among sexually active women and men in the U.S., it is reassuring to know that vaccines are now available, that these infections most often remain asymptomatic, that their symptoms, if they occur, are usually manageable, and that condom use is likely to reduce the risk of infection. Sexually active women should also be sure to have routine Pap and/or HPV tests as well.

Cited References

- ACOG – American College of Obstetricians and Gynecologists. (2000). "Make Decisions about Human Papillomavirus Based on Sound Medicine, Rather than Politics." _____ (2009, December). "Cervical Cytology Screening." *ACOG Practice Bulletin*, 109, 1-12.
- _____. (2013). "Management of Abnormal Cervical Cancer Screening Test Results and Cervical Cancer Precursors." *ACOG Practice Bulletin*, 140, 1-30.
- ACS – American Cancer Society. (2015a). *Cancer Facts and Figures 2015*. Atlanta, GA: American Cancer Society, Inc. [Online]. <http://www.cancer.org/acs/groups/content/@editorial/documents/document/acspc-044552.pdf>, accessed January 9, 2015.
- _____. (2015b). *Cervical Cancer*. Atlanta, GA: American Cancer Society, Inc. [Online]. <http://www.cancer.org/acs/groups/cid/documents/webcontent/003094-pdf.pdf>, accessed January 9, 2015.
- Anic, Gabriella M. and Anna R. Giuliano. (2012). "Genital HPV infection and related lesions in men." *Preventive Medicine*, 53 (Supp 1): S36-S41
- Anttila, Tarja, et al. (2001). "Serotypes of Chlamydia trachomatis and Risk for Development of Cervical Squamous Cell Carcinoma." *JAMA*, 285(1), 47-51.
- ASHA – American Social Health Association. (2014) *Genital Warts*. [Online]. <http://www.ashasexualhealth.org/stdsstis/hpv/genital-warts/>, accessed December 2, 2014.
- Bleeker, Maaike C.G., et al. (2003). "Condom Use Promotes Regression of Human Papillomavirus-Associated Penile Lesions in Male Sexual Partners of Women with Cervical Intraepithelial Neoplasia." *International Journal of Cancer*, 107, 804-10.
- Boehner, Constance W., et al. (2003). "Viral Sexually Transmitted Disease Vaccine Acceptability Among College Students." *Sexually Transmitted Diseases*, 30(10), 774-8.
- Boonstra, Heather. (2004). "Comprehensive Approach Needed to Combat Sexually Transmitted Infections Among Youth." *The Guttmacher Report on Public Policy*, 7(1), 3-4, 13.
- Bor, Jonathan. (2006, May 11). "HPV Vaccine Nearing Approval; FDA Likely to OK Immunizer That May Eliminate Most Cases of Cervical Cancer." *Baltimore Sun*, p. 1A.
- Bosch, F. Xavier, and Silvia de Sanjosé. (2003). "Chapter 1: Human Papillomavirus and Cervical Cancer – Burden and Assessment of Causality." *Journal of the National Cancer Institute Monograph*, 31, 3-13.
- Brown, Darron R., et al. (2005). "A Longitudinal Study of Genital Human Papillomavirus Infection in a Cohort of Closely Followed Adolescent Women." *Journal of Infectious Diseases*, 191, 182-92.
- Carson, Sylvia. (1997). "Human Papillomatous Virus Infection Update: Impact on Women's Health." *The Nurse Practitioner*, 22(4), 24-37.
- CDC – Centers for Disease Control and Prevention. (2012a). *HPV-Associated Cancers Diagnosis by Age*. [Online]. <http://www.cdc.gov/cancer/hpv/statistics/age.htm>, accessed December 3, 2014.
- _____. (2012b). *HPV Vaccine Information for Young Women – Fact Sheet*. Online, www.cdc.gov/std/hpv/stdfact-hpv-vaccine-young-women.htm, accessed December 3, 2014.
- _____. (2013). *Ready-To-Use STD Curriculum - Human Papillomavirus (HPV) Module - Faculty Notes*. [Online]. <http://www2a.cdc.gov/stdtraining/ready-to-use/Manuals/HPV/hpv-notes-2013.pdf>, accessed December 2, 2014.
- _____. (2014a). *Genital HPV Infection - Fact Sheet*. [Online]. <http://www.cdc.gov/std/HPV/STDFact-HPV.htm>, accessed 2014, December 2.
- _____. (2014b). *HPV Vaccine - Questions & Answers*. <http://www.cdc.gov/vaccines/vpd-vac/hpv/vac-faqs.htm>, accessed December 2, 2014.
- _____. (2014c). *Human Papillomavirus (HPV): Signs and Symptoms*. [Online]. <http://www.cdc.gov/hpv/signs-symptoms.html>, accessed December 2, 2014.
- "The Cervical Cancer Virus." (1995). *Discover*, 16, 24-6.
- Charturvedi, Anil K. et al. (2008). "Incidence Trends for Human Papillomavirus-Related and -Unrelated Oral Squamous Cell Carcinomas in the United States." *Journal of Clinical Oncology*, 26(4), 612-619.
- Clifford, G.M., et al. (2003). "Human Papillomavirus Types in Invasive Cervical Cancer Worldwide: A Meta-Analysis." *British Journal of Cancer*, 88(1), 63-73.
- Cockerell, Clay J. (1995). "Human Papillomavirus Infections. International Association of Physicians in AIDS Care." [Online]. <http://www.iapac.org/clinmgmt/diseases/cm/cm6.html#1>, accessed October 1, 1999.
- Davis, Kristin, et al. (2004). "Human Papillomavirus Vaccine Acceptability Among Parents of 10- to 15- Year Old Adolescents." *Journal of Lower Genital Tract Disease*, 8(3), 188-94.
- Dunne, Eileen F., et al. (2007). "Prevalence of HPV Infection among Females in the United States." *JAMA*, 297(8), 813-819.
- Elfgren, Kristina, et al. (2000). "A Population-Based Five-Year Follow-Up Study of Cervical Human Papillomavirus Infection." *American Journal of Obstetrics and Gynecology*, 183(3), 561-7.
- Eng, Thomas, and William Butler, eds. (1997). *The Hidden Epidemic: Confronting Sexually Transmitted Diseases*. Washington, DC: National Academy Press.

- FDA – Food and Drug Administration. (2006, June 8). *FDA News: FDA Licenses New Vaccine for Prevention of Cervical Cancer and Other Diseases in Females Caused by Human Papillomavirus*. [Online]. <http://www.fda.gov/bbs/topics/NEWS/2006/NEW01385.html/>, accessed July 12, 2006.
- _____. (2014, December 10). "FDA approves Gardasil 9 for prevention of certain cancers caused by five additional types of HPV." Press release. [Online]. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm426485.htm>, accessed December 16, 2014.
- Ferlay J., et al. (2014). *GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11*. Lyon, France: International Agency for Research on Cancer. [Online]. http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx?cancer=cervix, accessed December 3, 2014. [Internet].
- Friedman, Allison L. and Hilda Sheppard. (2007). "Exploring the Knowledge, Attitudes, Beliefs, and Communication Preferences of the General Public Regarding HPV: Findings from CDC Focus Group Research and Implications for Practice." *Health Education and Behavior*, 34 (3), 471-485.
- Gilden, David. (2005). "Protecting Against HPV." *GMHC Treatment Issues*, 19(5/6). [Online]. <http://www.gmhc.org/health/treatment/ti/ti1956.html>, accessed August 10, 2006.
- Gillison, Maura L. et al. (2012). "Prevalence of Oral HPV in the United States, 2009-2010." *JAMA*, 307(7), 693-703.
- GlaxoSmithKline (GSK). (2006). New Data Show Cervarix™, GSK's HPV 16/18 Cervical Cancer Candidate Vaccine, Is Highly Immunogenic and Well-Tolerated in Women Over 25 Years of Age. [Online]. http://www.gsk.ca/en/media_room/news/20060605.pdf, accessed July 7, 2006.
- González Intxaurreaga, M.A., et al. (2002). "HPV and Carcinogenesis." *Acta Dermatovenerol*, 11(3), 1-8.
- GSK – GlaxoSmithKline. (2009). "FDA Approves Cevarix, GlaxoSmithKline's Cancer Vaccine." Press release, October 16.
- Harper, Diane M., et al. (2004). "Efficacy of a Bivalent L1 Virus-Like Particle Vaccine in Prevention of Infection with Human Papillomavirus Types 16 and 18 in Young Women: A Randomised Controlled Trial." *Lancet*, 364, 1757-65.
- _____. (2006). "Sustained Efficacy Up to 4.5 Years of a Bivalent L1 Virus-Like Particle Vaccine Against Human Papillomavirus Types 16 and 18: Follow-Up From a Randomised Control Trial." *Lancet*. DOI: 10.1016/S0140-6736(06)68439-0.
- Hildesheim, A., et al. (1994). "Persistence of Type-Specific Human Papillomavirus Infection Among Cytologically Normal Women." *Journal of Infectious Diseases*, 169(2), 235-40.
- Ho, Gloria Y.F., et al. (1995). "Persistent Genital Human Papillomavirus Infection as a Risk Factor for Persistent Cervical Dysplasia." *Journal of the National Cancer Institute*, 87(18), 1365-71.
- Ho, Gloria Y.F., et al. (1998). "Natural History of Cervicovaginal Papillomavirus Infection in Young Women." *New England Journal of Medicine*, 338(7), 423-8.
- Hogewoning, Cornelis J.A., et al. (2003). "Condom Use Promotes Regression of Cervical Intraepithelial Neoplasia and Clearance of Human Papillomavirus: A Randomized Trial." *International Journal of Cancer*, 107, 811-6.
- Holmes, King, et al, eds. (2008). *Sexually Transmitted Diseases*, Fourth Ed. New York: McGraw-Hill Medical.
- "HPV DNA Tests: Studies Target Use for Cancer Screening." (2000, May). *STD Quarterly*, 1-3.
- Insinga, Ralph P., et al. (2004). "The Health Care Costs of Cervical Human Papillomavirus-Related Disease." *American Journal of Obstetrics & Gynecology*, 191, 114-20.
- IOM – Institute of Medicine. (2011, August). "Adverse Effects of Vaccines: Evidence and Causality." Washington, DC: IOM.
- Janicek, Mike, and Hervy Averette. (2001). "Cervical Cancer: Prevention, Diagnosis, and Therapeutics." *CA: A Cancer Journal for Clinicians*, 51(2), 92-114.
- Jay, Naomi, and Anna-Barbara Moscicki. (2000). "Human Papilloma Virus Infection in Women." In Marlene Goldman & Maureen Hatch, eds., *Women and Health*. San Diego, CA: Academic Press.
- Kahn, Jessica A. (2005). "Vaccination As A Prevention Strategy for Human Papillomavirus-Related Diseases." *Journal of Adolescent Health*, 37, S10-6.
- Kahn, Jessica A., et al. (2003). "Attitudes About Human Papillomavirus Vaccine in Young Women." *International Journal of STD & AIDS*, 14, 300-6.
- Kahn, Jessica A., et al. (2005). "Pediatricians' Intention to Administer Human Papillomavirus Vaccine: The Role of Practice Characteristics, Knowledge, and Attitudes." *Journal of Adolescent Health*, 37, 502-10.
- Kann, Laura, et al. (2014). "Youth Risk Behavior Surveillance – United States, 2013." *Morbidity and Mortality Weekly Report*, 63 (SS-4), 1-96. [Online]. <http://www.cdc.gov/mmwr/pdf/ss/ss6304.pdf>, accessed December 3, 2014.
- Kashima, H.K., et al. (1996). "Recurrent Respiratory Papillomatosis." *Obstetrics and Gynecologic Clinics of North America*, 23(3), 699-706.
- Kay, Jane. (2005, February 1, accessed 2006, May 19). "X-Rays Added To Cancer List; Some Viruses Also Among Carcinogens on Federal Registry." *San Francisco Chronicle*, p. A1.
- Keller, Mary L., et al. (1995). "Genital Human Papillomavirus Infection: Common But Not Trivial." *Health Care for Women International*, 16, 351-64.
- Kjaer, Susanne Krüger, et al. (2001). "High-Risk Human Papillomavirus Is Sexually Transmitted: Evidence From a Follow-Up Study of Virgins Starting Sexual Activity (Intercourse)." *Cancer Epidemiology, Biomarkers & Prevention*, 10(2), 101-6.
- Koutsky, Laura, and Nancy Kiviat. (1999). "Genital Human Papillomavirus." In King Holmes, et al., eds., *Sexually Transmitted Diseases*, 3rd ed. New York: McGraw-Hill.

- Koutsky, Laura A., et al. (2002). "A Controlled Trial of a Human Papillomavirus Type 16 Vaccine." *New England Journal of Medicine*, 347(21), 1645-51.
- Kreimer, Aimée R. et al. (2011). "Efficacy of a bivalent HPV 16/18 vaccine against anal HPV 16/18 infection among young women: a nested analysis within the Costa Rica Vaccine Trial." *The Lancet Oncology*. DOI: 10.1016/S1473-0245(11)70213-3.
- Lytwyn, Alice, and John W. Sellors. (1997). "Sexually Transmitted Human Papillomaviruses: Current Concepts and Control Issues." *The Canadian Journal of Human Sexuality*, 6(2), 113-26.
- Mao, Constance, et al. (2006). "Efficacy of Human Papillomavirus-16 Vaccine to Prevent Cervical Intraepithelial Neoplasia: A Randomized Controlled Trial." *Obstetrics & Gynecology*, 107(1), 18-27.
- Markowitz, Lauri E., et al. (2014, August 29). "Human Papillomavirus Vaccination: Recommendations of the Advisory Committee on Immunization Practices (ACIP)." *Morbidity and Mortality Weekly Report*, 63(RR05): 1-30. [Online]. <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6305a1.htm>, accessed December 3, 2014.
- Mays, Rose M., et al. (2004). "Parental Perspectives on Vaccinating Children Against Sexually Transmitted Infections." *Social Science & Medicine*, 58, 1405-13.
- McDermott-Webster, Marian. (1999). "The HPV Epidemic." *American Journal of Nursing*, 99, 24L-24N.
- Merck. (2014, December 11). "FDA Approves Merck's HPV Vaccine, GARDASIL®9, to Prevent Cancers and Other Diseases Caused by Nine HPV types - Including Types that Cause About 90% of Cervical Cancer Cases." Press release. [Online]. <http://www.mercknewsroom.com/news-release/prescription-medicine-news/fda-approves-mercks-hpv-vaccine-gardasil9-prevent-cancers-an>, accessed December 16, 2014
- Moscicki, Anna-Barbara. (2005). "Impact of HPV Infection in Adolescent Populations." *Journal of Adolescent Health*, 37, S3-9.
- Moscicki, Anna-Barbara, et al. (1998). "The Natural History of Human Papillomavirus Infection As Measured By Repeated DNA Testing in Adolescent and Young Women." *Journal of Pediatrics*, 132(2), 277-84.
- Muñoz, Nubia, et al. (2003). "Epidemiologic Classification of Human Papillomavirus Types Associated with Cervical Cancer." *New England Journal of Medicine*, 348(6), 518-27.
- Muñoz, Nubia, et al. (2004). "Incidence, Duration, and Determinants of Cervical Human Papillomavirus Infection in a Cohort of Colombian Women with Normal Cytological Results." *Journal of Infectious Diseases*, 190, 2077-87.
- _____. (2010). "Impact of Human Papillomavirus (HPV)-6/11/16/18 Vaccine on All HPV-Associated Genital Diseases in Young Women." *JNCI*, 102 (5), 325-339.
- NCCC - National Cervical Cancer Coalition. (2014). *Screening: Pap and HPV tests*. [Online]. <http://www.nccc-online.org/index.php/screening>, accessed December 16, 2014.
- NCI - National Cancer Institute. (2014). *Cervical Cancer Prevention (PDQ®)*. [Online]. <http://www.cancer.gov/cancertopics/pdq/prevention/cervical/HealthProfessional>, accessed December 2, 2014.
- _____. (2011). *NCI Fact Sheet - Human Papillomavirus (HPV) Vaccines*. Online, www.cancer.gov/cancertopics/factsheet/prevention/HPV-vaccine, accessed November 5, 2014
- _____. (2013). "HINTS - Health Information National Trends Survey." Online. <http://hints.cancer.gov/question-details.aspx?qid=526&dataset=43&method=mail>, accessed November 5, 2014.
- NIAID - National Institute of Allergy and Infectious Diseases. National Institutes of Health. Public Health Service. (2004). *Human Papillomavirus and Genital Warts*. [Online]. <http://www.niaid.nih.gov/factsheets/stdhvp.htm>, accessed May 23, 2006.
- Owusu-Edusei, Kwame Jr., et al. (2013). "The Estimated Direct Medical Cost of Selected Sexually Transmitted Diseases in the United States, 2008." *Sexually Transmitted Diseases*, 40(3), 197-201.
- Puranen, Mirja, et al. (1997). "Exposure of an Infant to Cervical Human Papillomavirus Infection of the Mother is Common." *American Journal of Obstetrics and Gynecology*, 176(5), 1039-45.
- Richardson, Harriet, et al. (2003). "The Natural History of Type-Specific Human Papillomavirus Infections in Female University Students." *Cancer Epidemiology, Biomarkers & Prevention*, 12, 485-90.
- Rosenthal, Susan L., et al. (1995). "Hepatitis B Vaccine Acceptance among Adolescents and Their Parents." *Journal of Adolescent Health*, 17, 248-54.
- Ryerson, A.B., et al. (2008). "Burden of potentially human papillomavirus-associated cancers of the oropharynx and oral cavity in the US, 1998-2003." *Cancer*, 113 (10 Suppl), 2901-9.
- Schiffman, Mark, and Philip E. Castle. (2003). "Human Papillomavirus: Epidemiology and Public Health." *Archives of Pathology and Laboratory Medicine*, 127, 930-4.
- Schwarz, T.F., et al. (2006). "An AS04-containing human papillomavirus (HPV) 16/18 vaccine for prevention of cervical cancer is immunogenic and well-tolerated in women 15-55 years old." *Journal of Clinical Oncology*, 24(18S), abstract 1008.
- Serrano, Beatriz, et al. (2014). "Potential impact of a 9-valent HPV vaccine in HPV-related cervical disease in 4 emerging countries (Brazil, Mexico, India and China)." *Cancer Epidemiology*, 38(6), 748-756.
- Shah, Keerti V. (1997). "Human Papillomaviruses and Anogenital Cancers." *The New England Journal of Medicine*, 337(19), 1386-8.

- Skjeldestad, Finn Egil, et al. (2005). "Prophylactic Quadrivalent Human Papillomavirus (HPV) (Types 6, 11, 16, 18) L1 Virus-Like Particle (VLP) Vaccine (Gardasil™) Reduced Cervical Intraepithelial Neoplasia (CIN) 2/3 Risk." Presented at: Infectious Disease Society of America 43rd Annual Meeting; October 7, 2005; San Francisco, California. Abstract LB-8a. [Online]. <http://www.idsociety.org/Template.cfm?Section=Program2&CONTENTID=14108&TEMPLATE=/ContentManagement/ContentDisplay.cfm>, accessed May 26, 2006.
- Smith, Elaine M., et al. (2004). "Human Papillomavirus Prevalence and Types in Newborns and Parents: Concordance and Modes of Transmission." *Sexually Transmitted Diseases*, 31(1), 57-62.
- Soper, David E. (2002). "Genitourinary Infections and Sexually Transmitted Diseases." In Jonathan S. Berek, ed., *Novak's Gynecology*, 13th ed. Philadelphia, PA: Lippincott Williams & Williams.
- Stevens-Simon, Catherine, et al. (2000). "The Prevalence of Genital Human Papillomavirus Infections in Abused and Nonabused Preadolescent Girls." *Pediatrics*, 106(4), 645-9.
- Stone, Katherine, et al. (1999). "Barrier Methods for the Prevention of Sexually Transmitted Diseases." In King Holmes, et al., eds., *Sexually Transmitted Diseases*, 3rd ed. New York: McGraw-Hill.
- Taira, Al V., et al. (2004). "Evaluating Human Papillomavirus Vaccination Programs." *Emerging Infectious Diseases*, 10(11), 1915-23.
- Tuller, David. (2003, February 18). "Some Urge Type of Pap Test To Find Cancer in Gay Men." *The New York Times*, p. F7.
- Verdon, Mary. (1997). "Issues in the Management of Human Papillomavirus Genital Disease." *American Family Physician*, 55, 1813-20.
- Villa, Luisa L., et al. (2005). "Prophylactic Quadrivalent Human Papillomavirus (Types 6, 11, 16, and 18) L1 Virus-Like Particle Vaccine in Young Women: A Randomised Double-Blind Placebo-Controlled Multicentre Phase II Efficacy Trial." *Lancet Oncology*, 6, 271-8.
- Walboomers, J.M., et al. (1999). "Human Papillomavirus Is a Necessary Cause of Invasive Cervical Cancer Worldwide." *Journal of Pathology*, 189(1), 12-9.
- Weinstock, Hillard, et al. (2004). "Sexually Transmitted Diseases Among American Youth: Incidence and Prevalence Estimates, 2000." *Perspectives on Sexual and Reproductive Health*, 36(1), 6-10.
- Wiley, D.J., et al. (2002). "External Genital Warts: Diagnosis, Treatment, and Prevention." *Clinical Infectious Diseases*, 35(Suppl 2), S210-24.
- Winer, Rachel L., et al. (2003). "Genital Human Papillomavirus Infection: Incidence and Risk Factors in a Cohort of Female University Students." *American Journal of Epidemiology*, 157(3), 218-26.
- Winer, Rachel L., et al. (2006). "Condom Use and the Risk of Genital Human Papillomavirus Infection of Young Women." *New England Journal of Medicine*, 354(25), 2645-54.
- Woodman, Ciaran B.J., et al. (2001). "Natural History of Cervical Human Papillomavirus Infection In Young Women: A Longitudinal Cohort Study." *Lancet*, 357, 1831-6.

Media Contact – 212-261-4433
Last updated June 2015