

# THE DIFFERENCE BETWEEN THE MORNING-AFTER PILL AND THE ABORTION PILL

There has been considerable public confusion about the difference between the morning-after pill and the abortion pill because of misinformation disseminated by groups that oppose safe and legal abortion. The morning-after pill, also known as emergency contraception, helps *prevent* pregnancy; the abortion pill, also known as medication abortion, *terminates* pregnancy. According to the general medical definitions of pregnancy that have been endorsed by many organizations — including the American College of Obstetricians and Gynecologists and the United States Department of Health and Human Services — pregnancy begins when a pre-embryo completes implantation into the lining of the uterus (ACOG, 1998; DHHS, 1978; Hughes, 1972; “Make the Distinction ...,” 2001). Hormonal methods of contraception, including the morning-after pill, prevent pregnancy by inhibiting ovulation and fertilization (ACOG, 1998). The abortion pill terminates a pregnancy without using instruments. By helping women terminate unwanted pregnancies up to 63 days after their last menstruation, the abortion pill is a safe and effective option.

## THE MORNING-AFTER PILL

### What is the morning after pill?

Also known as emergency contraception, the morning-after pill contains medication that reduces the risk of pregnancy if started within 120 hours (five days) of unprotected intercourse. Plan B One-Step® contains the hormone progestin and are currently available over the counter in the family planning aisle of drug stores with no age requirement. Other brands of levonorgestrel

EC such as Next Choice® remain behind the counter with pharmacists for purchase by anyone 17 or older without a prescription, or anyone younger than 17 with a prescription. ella®, which contains ulipristal acetate (UPA), and certain brands of oral contraception taken in increased doses for use as emergency contraception require a prescription at any age (Barr Pharmaceuticals, 2006; Glasier, 2010; RHTP, 2009; Rodrigues et al., 2001; Van Look & Stewart, 1998).

## THE ABORTION PILL

### What is the abortion pill?

Also known as medication abortion, the abortion pill contains medication called mifepristone to induce abortion. Mifepristone (Mifeprex®) can be taken under supervision up to 63 days after the first day of the last menstrual period. It is used in conjunction with misoprostol, which is taken later to complete the abortion (Creinin & Aubény, 1999; Middleton et al., 2005; Schaff et al., 2000; Schaff et al., 2001).

## THE MORNING-AFTER PILL

### How does the morning-after pill work?

In its approval of the morning-after pill, the U.S. Food and Drug Administration (FDA) declared, “Emergency contraceptives act by delaying or inhibiting ovulation and/or altering tubal transport of sperm and/or ova (thereby inhibiting implantation)” (FDA, 1997). More recently, studies suggested that progestin-only morning-after pills work only by preventing ovulation or fertilization, and have no effect on implantation (Croxatto et al., 2003; Novikova et al., 2007). In 2008, a consortium of authorities declared that progestin-only emergency contraception does not interfere with implantation (ICEC-FIGO, 2008). UPA works only by preventing ovulation (Glasier, 2010).

### How effective is the morning-after pill?

The morning-after pill is very effective at reducing the risk of pregnancy. Studies have shown that it reduces the risk of pregnancy when taken up to 120 hours after unprotected intercourse. With the exception of UPA, the sooner the dosing begins, the more effective the treatment. When taken within 72 hours of unprotected intercourse, morning-after pills that contain both estrogen and progestin reduce the risk of pregnancy by 75 percent. Within that same time, progestin-only regimens, such as Plan B One-Step and Next Choice, reduce the risk of pregnancy by 89 percent. When initiated within 24 hours of unprotected intercourse, progestin-only morning-after pill reduced the risk of pregnancy by 95 percent (Ellertson et al., 2003; Rodrigues et al., 2001; TFPMFR, 1998; Van Look & Stewart, 1998). The effectiveness of UPA, however, does not diminish over the course of the five days following unprotected intercourse (Fine et al., 2010; Glasier et al., 2010).

## THE ABORTION PILL

### How does the abortion pill work?

Mifepristone ends pregnancy by blocking the hormones necessary for maintaining a pregnancy. Misoprostol causes the uterus to contract and empty (Creinin & Aubény, 1999).

### How effective is the abortion pill?

The abortion pill is highly effective at ending very early pregnancies. Complete abortion will occur in 96–97 percent of women who choose mifepristone. In the small percentage of cases that medication abortion fails, other abortion procedures are required to end the pregnancies (ACOG, 2001; Schaff et al., 2000).

## THE MORNING-AFTER PILL

### How safe is the morning-after pill?

The morning-after pill is safe for nearly all women — millions of women around the world have used it safely (Guillebaud, 1998; Van Look & Stewart, 1998).

### Does the morning-after pill cause an abortion?

The morning-after pill will not induce an abortion in a woman who is already pregnant, nor will it affect the developing pre-embryo or embryo (Van Look & Stewart, 1998). Emergency contraception prevents pregnancy and helps a woman prevent the need for abortion.

### Why might a woman choose the morning-after pill?

Women may choose emergency contraception as a way to prevent pregnancy after unprotected intercourse — in cases of unanticipated sexual activity, contraceptive failure, or sexual assault. Nearly half of America's 6.7 million annual pregnancies are unintended (Finer & Zolna, 2011).

### Does the morning-after pill have side effects?

Side effects are far less common using progestin-only and UPA pills than using combined hormone pills. The most common side effects include nausea and vomiting. Abdominal pain, breast tenderness, dizziness, fatigue, headaches, and irregular bleeding may also occur (Van Look & Stewart, 1998; OPR, 2011; TFPMFR, 1998; Trussell & Schwarz, 2011).

## THE ABORTION PILL

### How safe is the abortion pill?

The abortion pill is safe for most women — millions of women around the world have used it safely. There are risks associated with all medical procedures, including abortion. And, in extremely rare cases, death is possible from serious complications of the abortion pill, but it remains safer than carrying a pregnancy to term (ARHP, 2008).

### Can the medicines used in the abortion pill also be used for emergency contraception?

Although some studies show that mifepristone could be used in very low doses to reduce the risk of pregnancy as a method of emergency contraception within five days of unprotected intercourse, it is not approved for use as emergency contraception in the United States at this time (Ho et al., 2002; TFPMFR, 1999).

### Why might a woman choose the abortion pill?

Women might choose the abortion pill as a way to end pregnancy because it is a noninvasive procedure and does not require anesthesia. It is free from the risk of injury to the cervix or uterus and the possible complications caused by the use of anesthesia used for other abortion procedures (Aguillaume & Tyrer, 1995). Women who chose medication abortion also reported that they felt it was a more “natural” way to end a pregnancy (Winikoff, 1995).

### Does the abortion pill have side effects?

The most common side effects following medication abortion are similar to those of a miscarriage — abdominal pain, bleeding, changes in body temperature, dizziness, fatigue, and gastrointestinal distress (ACOG, 2005; Creinin & Aubény, 1999; Stewart et al., 2005).

## THE MORNING-AFTER PILL

### How long does the process of using the morning-after pill take?

Combined hormone pills are taken in two doses, 12 hours apart. Progestin-only pills can be taken in one dose. UPA is taken in one dose. Side effects associated with emergency contraception pills generally subside within 48 hours. They affect the timing of the menstrual cycle in 10–15 percent of women. Changes in the menstrual cycle are seen with combined hormone, progestin-only, and UPA pills. If the next menstrual cycle is more than one week late, a woman should visit her clinician for a pregnancy test (Fine et al., 2010; Van Look & Stewart, 1998; von Hertzen et al., 2002).

### Are women who have used the morning-after pill satisfied with it?

An overwhelming majority of morning-after pill users are satisfied with it. One study found that 97 percent of users would recommend it to friends and family (Harvey et al., 1999). Another study found that 92 percent of women who had used emergency contraception would use it again in the case of a contraceptive emergency (Breitbart et al., 1998).

### Where can I get the morning-after pill?

Plan B-One Step is now available over the counter in the family planning aisle of drug stores with no age requirement. Other brands of levonorgestrel

EC remain behind the counter with pharmacists for purchase by anyone 17 or older without a prescription, or anyone younger than 17 with a prescription. ella requires a prescription at any age. If you need a prescription for emergency contraception, you can contact your nearest Planned Parenthood health center at 1-800-230-PLAN or [www.plannedparenthood.org](http://www.plannedparenthood.org).

## THE ABORTION PILL

### How long does the process of the abortion pill take?

It begins immediately after taking the mifepristone. Some women may begin spotting *before* taking the misoprostol, the second medication. For most, the bleeding and cramping associated with medication abortion begin *after* taking it. More than 50 percent of women who use mifepristone abort within four or five hours after taking the misoprostol. Heavy bleeding may continue for about 13 days. Spotting can last for a few weeks. About 92 percent of mifepristone abortions are completed within a week (ACOG, 2001; el-Refaey et al., 1995; Newhall & Winikoff, 2000; Peyron et al., 1993; Wiebe et al., 2002).

### Are women who have used the abortion pill satisfied with the method?

An overwhelming majority of women who choose medication abortion were satisfied with it. A recent study found that 97 percent of women who had a medication abortion would recommend it to a friend. Additionally, 91 percent of the women reported that they would choose medication abortion again if they had to have another abortion (Hollander, 2000).

### Where can I get the abortion pill?

Contact your nearest Planned Parenthood health center at 1-800-230-PLAN or [www.plannedparenthood.org](http://www.plannedparenthood.org), another women's health care center, or your private clinician. Planned Parenthood health centers that do not provide medication abortion can refer you to a provider who does.

### How much does the morning-after pill cost?

Nationwide, the price of EC ranges from \$30–\$65 (PPFA, 2013b). Costs vary from community to community, based on regional and local expenses. Contact your nearest Planned Parenthood health center at 1-800-230-PLAN or [www.plannedparenthood.org](http://www.plannedparenthood.org), another women's health care center, or your private clinician.

### How much does the abortion pill cost?

Nationwide, the price of medication abortion ranges between \$300 and \$800. This includes two or three office visits, testing, and exams (PPFA, 2013a). Costs vary from community to community, based on regional and local expenses.

### CITED REFERENCES

- ACOG — American College of Obstetricians and Gynecologists. (1998, July). Statement on Contraceptive Methods.
- \_\_\_\_\_. (2001, April). "Medical Management of Abortion." *ACOG Practice Bulletin*, 26, 1–13.
- \_\_\_\_\_. (2005, October). "Medical Management of Abortion." *ACOG Practice Bulletin*, 67, 1–12.
- Aguillaume, Claude & Louise Tyrer. (1995). "Current Status and Future Projections on Use of RU-486." *Contemporary Ob/Gyn*, 40(6), 23–40.
- ARHP — Association of Reproductive Health professionals. (2008, April). *What You Need to Know — Mifepristone Safety Overview*. [Online]. <http://www.arhp.org/upload/Docs/mifepristonefactsheet.pdf>, accessed September 29, 2009.
- Barr Pharmaceuticals, Inc. (2006, August 24). *FDA Grants OTC Status to Barr's Plan B® Emergency Contraceptive: Historic Dual Status Decision Provides OTC Access to Those 18 Years of Age and Older; Remains Prescription for Women 17 and Younger*. [Online]. <http://phx.corporate-ir.net/phoenix.zhtml?c=irol-newsArticle&ID=899120>.
- Breitbart, Vicki, et al. (1998). "The Impact of Patient Experience on Practice: The Acceptability of Emergency Contraceptive Pills in Inner City Clinics." *Journal of the American Medical Women's Association*, 53(5 Supplement 2), 255–58.
- Creinin, Mitchell & Elizabeth Aubény. (1999). "Medical Abortion in Early Pregnancy." In Maureen Paul, et al., Eds. *A Clinician's Guide to Medical and Surgical Abortion*. New York: Churchill Livingstone.
- Croxatto, Horatio B., et al. (2003). "Mechanisms of Action of Emergency Contraception." *Steroids*, 68, 1095–98.
- DHHS — U.S. Department of Health and Human Services. (1978). *Code of Federal Regulations*. 45CFR46.203.
- Ellertson, Charlotte, et al. (2003). "Extending the Time Limit for Starting the Yuzpe Regimen of Emergency Contraception to 120 hours." *Obstetrics and Gynecology*, 101, 1168–71.
- El-Refaey, H., et al. (1995). "Induction of Abortion with Mifepristone (RU 486) and Oral or Vaginal Misoprostol." *New England Journal of Medicine*, 332(15), 983–7.
- FDA — U.S. Food and Drug Administration. (1997). "Prescription Drug Products; Certain Combined Oral Contraceptives for Use as Postcoital Emergency Contraception." *Federal Register*, 62(37), 8609–12.
- Fine, Paul T. et al. (2010). "Ulipristal Acetate Taken 48–120 Hours after Intercourse for Emergency Contraception." *Obstetrics and Gynecology*, 115(2), 1–7.
- Finer, Lawrence B. & Mia R. Zolna. (2011). "Unintended pregnancy in the United States: incidence and disparities, 2006." *Contraception*, 84(5), 478–485.
- Glasier, Anne F. et al. (2010). "Ulipristal acetate versus levonorgestrel for emergency contraception: a randomized non-inferiority trial and meta-analysis." *The Lancet*, 365, 555–62.
- Guillebaud, John. (1998). "Commentary: Time for Emergency Contraception with Levonorgestrel Alone." *The Lancet*, 352(9126), 416.
- Harvey, S. Marie, et al. (1999). "Women's Experience and Satisfaction with Emergency Contraception." *Family Planning Perspectives*, 31(5), 237–40 & 260.
- Ho, Park Chung, et al. (2002). "Mifepristone: Contraceptive and Non-Contraceptive Uses." *Current Opinions in Obstetrics and Gynecology*, 14(3), 325–30.
- Hollander, Dore. (2000). "Most Abortion Patients View Their Experience Favorably, But Medical Abortion Gets a Higher Rating Than Surgical." *Family Planning Perspectives*, 32(5), 264.
- Hughes, Edward, Ed. (1972). *Obstetric-Gynecologic Terminology*. Philadelphia, PA: F. A. Davis Company.

- ICEC-FIGO — International Consortium for Emergency Contraception – International Federation of Gynecology & Obstetrics. (2008-October). “How do levonorgestrel-only emergency contraceptive pills (LNG ECPs) prevent pregnancy?” [Online]. [http://www.cecinfo.org/PDF/ICEC\\_MOA\\_10\\_14.pdf](http://www.cecinfo.org/PDF/ICEC_MOA_10_14.pdf), accessed September 25, 2009.
- “Make the Distinction: EC Prevents Pregnancy.” (2001). *Contraceptive Technology Update*, 22(1),4.
- Middleton, Tamer, et al., (2005). “Randomized Trial of Mifepristone and Buccal or Vaginal Misoprostol for Abortion Through 56 Days of Last Menstrual Period.” *Contraception*, 72, 328–32.
- Newhall, Elizabeth Pirruccello & Beverly Winikoff, (2000). “Abortion with Mifepristone and Misoprostol: Regimens, Efficacy, Acceptability and Future Directions.” *American Journal of Obstetrics and Gynecology*, 183(2), S44–53.
- Novikova, Natalia, et al. (2007). “Effectiveness of levonorgestrel emergency contraception given before or after ovulation — a pilot study.” *Contraception*, 75, 112–118.
- OPR — Office of Population Research, Princeton University. (2011, February 22, accessed April 21, 2011). “Answers to Frequently Asked Questions About... Types of Emergency Contraception.” [Online]. <http://ec.princeton.edu/questions/dose.html>, <http://ec.princeton.edu/questions/eceffect.html>, and <http://ec.princeton.edu/questions/ecsideeffects>.
- Peyron, R., et al. (1993). “Early Termination of Pregnancy with Mifepristone (RU 486) and Orally Active Prostaglandin Misoprostol.” *New England Journal of Medicine*, 328(21), 1509–13.
- PPFA — Planned Parenthood Federation of America. (2013a). “The Abortion Pill (Medication Abortion).” [Online]. <http://www.plannedparenthood.org/health-topics/abortion/abortion-pill-medication-abortion-4354.asp>, accessed December 5, 2013.
- \_\_\_\_\_. (2013b). “Emergency Contraception (Morning After Pill).” [Online]. <http://plannedparenthood.org/health-topics/emergency-contraception-morning-after-pill-4363.asp>, accessed December 5, 2013.
- RHTP — Reproductive Health Technologies Project. (2009). “FDA Approved Emergency Contraceptive Products Currently on the U.S. Market.” [Online]. [http://www.rhtp.org/contraception/emergency/documents/FDAApprovedEmergencyContraceptiveChartDecember2011-PRINTABLE\\_000.pdf](http://www.rhtp.org/contraception/emergency/documents/FDAApprovedEmergencyContraceptiveChartDecember2011-PRINTABLE_000.pdf), accessed January 10, 2012.
- Rodrigues, Isabel, et al. (2001). Effectiveness of Emergency Contraceptive Pills Between 72 and 120 Hours After Unprotected Sexual Intercourse.” *American Journal of Obstetrics and Gynecology*, 184(4), 416.
- Schaff, Eric, et al. (2000). “Low-Dose Mifepristone Followed by Vaginal Misoprostol at 48 Hours for Abortion up to 63 Days.” *Contraception*, 61(1), 41–6.
- Schaff, Eric, et al. (2001). “Randomized Trial of Oral Versus Vaginal Misoprostol at One Day after Mifepristone for Early Medical Abortion.” *Contraception*, 64, 81–5.
- Stewart, Felicia H., et al. (2005). “Abortion.” Pp. 673–700 in Robert A. Hatcher, et al., Eds., *Contraceptive Technology*— 18th Revised Edition. New York: Ardent Media, Inc.
- TFPMFR — Task Force on Postovulatory Methods of Fertility Regulation. (1998). “Randomised Controlled Trial of Levonorgestrel Versus the Yuzpe Regimen of Combined Oral Contraceptives for Emergency Contraception.” *The Lancet*, 352(9126), 428–33.
- \_\_\_\_\_. (1999). “Comparison of Three Single Doses of Mifepristone as Emergency Contraception: A Randomised Trial.” *The Lancet*, 353(9154), 697–702.
- Trussell, James, and Eleanor Bimla Schwarz. (2011). “Emergency Contraception.” Pp. 113–145 in Robert A. Hatcher et al., eds., *Contraceptive Technology* — 20th Revised Edition. New York: Ardent Media, Inc.
- Van Look, Paul & Felicia Stewart. (1998). “Emergency Contraception.” In Robert A. Hatcher et al., Eds, *Contraceptive Technology* — 17th Revised Edition. New York: Ardent Media.
- von Hertzen, Helena, et al. (2002). “Low Dose Mifepristone and Two Regimens of Levonorgestrel for Emergency Contraception: A WHO Multicentre Randomised Trial.” *The Lancet*, 260, 1803–10.
- Wiebe, Ellen, et al. (2002). “Comparison of Abortions Induced by Methotrexate or Mifepristone Followed by Misoprostol.” *Obstetrics and Gynecology*, 99(5), 813–9.
- Winikoff, Beverly. (1995). “Acceptability of Medical Abortion in Early Pregnancy.” *Family Planning Perspectives*, 27(4), 142–8, 185, & 199.