

**Why is this medication prescribed?**

Metronidazole eliminates bacteria and other microorganisms that cause infections of the reproductive system, gastrointestinal tract, skin, vagina, and other areas of the body. Antibiotics will not work for colds, flu, or other viral infections.

This medication is sometimes prescribed for other uses; ask your clinician or pharmacist for more information.

**How should this medicine be used?**

Metronidazole comes as a tablet to take by mouth. It is usually taken two or three times a day for 5-10 days or longer. Follow the directions on your prescription label carefully, and ask your clinician or pharmacist to explain any part you do not understand. Take metronidazole exactly as directed. Do not take more or less of it or take it more often than prescribed by your clinician.

**What special precautions should I follow?**

Before taking metronidazole,

- tell your clinician and pharmacist if you are allergic to metronidazole or any other drugs.
- tell your clinician and pharmacist what prescription and nonprescription medications you are taking, especially anticoagulants ('blood thinners') such as warfarin (Coumadin), astemizole (Hismanal), disulfiram (Antabuse), lithium (Lithobid), phenobarbital, phenytoin (Dilantin), and vitamins.
- tell your clinician if you have or have ever had blood, kidney, or liver disease or Crohn's disease.
- tell your clinician if you are pregnant, plan to become pregnant, or are breast-feeding. If you become pregnant while taking metronidazole, call your clinic.
- know that you should not drink alcohol while taking this drug. Alcohol may cause an upset stomach, vomiting, stomach cramps, headaches, sweating, and flushing (redness of the face).
- plan to avoid unnecessary or prolonged exposure to sunlight and to wear protective clothing, sunglasses, and sunscreen. Metronidazole may make your skin sensitive to sunlight.

**What should I do if I forget a dose?**

Take the missed dose as soon as you remember it. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. Do not take a double dose to make up for a missed one.

**What side effects can this medication cause?**

Metronidazole may cause side effects. Tell your clinician if any of these symptoms are severe or do not go away:

- vomiting
- diarrhea
- upset stomach
- loss of appetite
- stuffy nose
- dry mouth; sharp, unpleasant metallic taste
- dark or reddish-brown urine
- furry tongue; mouth or tongue irritation
- numbness or tingling of hands or feet

If you experience any of the following symptoms, call your clinic immediately:

- rash
- itching
- fever
- joint pain

If you experience a serious side effect, you or your clinician may send a report to the Food and Drug Administration's (FDA) MedWatch Adverse Event Reporting program online [at <http://www.fda.gov/MedWatch/report.htm>] or by phone [1-800-332-1088].

**What storage conditions are needed for this medicine?**

Keep this medication in the container it came in, tightly closed, and out of reach of children. Store it at room temperature and away from excess heat and moisture (not in the bathroom). Throw away any medication that is outdated or no longer needed. Talk to your pharmacist about the proper disposal of your medication.

**In case of emergency/overdose**

In case of overdose, call your local poison control center at 1-800-222-1222. If the victim has collapsed or is not breathing, call local emergency services at 911.

**What other information should I know?**

Keep all appointments with your clinic and the laboratory. Your clinician may order certain lab tests to check your response to metronidazole.

Do not let anyone else take your medication. Your prescription is probably not refillable. If you still have symptoms of infection after you finish the metronidazole, call your clinic.

- Flagyl®
- Flagyl® 375
- Flagyl® ER
- Helidac®  
Therapy

**Brand name(s):**

Last Revised –

04/01/2003 American Society of Health-System Pharmacists, Inc. Disclaimer - <http://www.nlm.nih.gov/medlineplus/ashpdisclaimer.html>



The MedMaster™ Patient Drug Information database provides information copyrighted by the American Society of Health-System Pharmacists, Inc., Bethesda, Maryland Copyright© 2004. All Rights Reserved.

**Note:** Licensed content is for personal use only. Do not forward this e-mail to mailing lists or distribute it for commercial purposes. See our copyright statement (<http://www.nlm.nih.gov/medlineplus/copyright.html>) for more information.